arXiv:1207.4898v1 [astro-ph.CO] 20 Jul 2012

Testing $f(R)$ dark energy model with the large scale structure

Jian-hua He[1]
[1] *INAF-Osservatorio Astronomico di Brera, Via Emilio Bianchi, 46, I-23807, Merate (LC), Italy*

In this work, we further investigate the family of $f(R)$ dark energy models that can exactly mimic the same background expansion history as that of the ΛCDM model. Instead of using the parameterized framework of modified gravity, we study the large scale structure in the $f(R)$ gravity using the full set cosmological perturbation equations. We investigate the structure formation in both the spatially flat and curved Universe. We also confront our model with the latest observations and conduct a Markov chain Monte Carlo analysis on the parameter space.

PACS numbers: 98.80.-k,04.50.Kd

I. INTRODUCTION

There has been accumulated conclusive evidence from supernovae[1] and other observations[2, 3] in the last decade, indicating that our Universe is undergoing a phase of accelerated expansion. Understanding the nature of the cosmic acceleration is one of the biggest questions in modern physics.

The leading explanation of the accelerated expansion is the cosmological constant within the context of General Relativity. However, the measured value of cosmological constant is far below the prediction of any sensible quantum field theories and the cosmological constant will inevitably lead to the coincidence problem that why the energy density of matter and vacuum are in the same order today(see [4] for review). Another possibility for the acceleration is that the universe is driven by a new and yet-unknown component called dark energy. The dark energy is some kind of dynamical fluid with negative and time-dependent equation of state $w(a)$. However, the nature of the dynamical dark energy is even harder to be understood in the fundamental physics than that of the cosmological constant.

Alternatively, a promising explanation of the acceleration is the modified gravity. The General relativity might not be ultimately correct in the cosmological scale. The Universe might be described by some kind of modified gravity. One simplest attempt is called $f(R)$ gravity, in which the scalar curvature in the Lagrange density of Einstein's gravity is replaced by an arbitrary function of R[5]. The $f(R)$ gravity can produce the accelerated expansion of the Universe with any designed effective dark energy equation of state w[6]. Furthermore, the time dependent effective DE EoS in the Jordan frame can be reproduced from the dilation of the inertial mass in the Einstein frame through the conformal transformation [7]. In this sense, the time dependent dark energy phenomenon can be better understood in a physical way in the framework of modified gravity.

In this paper, we investigate a specific family of $f(R)$ models that can reproduce the same background expansion history as that of the ΛCDM model since it has been argued that a valid $f(R)$ model should closely math the ΛCDM background[8]. The family of $f(R)$ model contains only one more extra parameter than that of the ΛCDM model and furthermore, it does have the well-defined Lagrangian formalism in the spatially flat Universe, which is valid for the whole expansion history of the Universe from the past to the future. The model is no longer simply a phenomenological model. Although this model has been studied by a number of work within the parameterized framework of modified gravity [9] [10], it still needs to solve the full set of cosmological perturbation equations to get more accurate results on the scale k dependent growth history of the Universe when confronted with upcoming high precision data in cosmological surveys. Therefore, in this work, instead of using the parameterized framework of modified gravity, we investigate the impact of $f(R)$ models on the large scale structure using the full set of covariant cosmological perturbation equations. We will confront our $f(R)$ model with the latest observations and conduct a Markov chain Monte Carlo analysis on the parameter space. We will also explore the spatially non-flat case in the $f(R)$ gravity, which has not been addressed in previous work [10] [11] [12].

This paper is organized as follows: In section II, we review the background dynamics of the Universe in the $f(R)$ gravity. In section III, we present the scalar perturbation equations in the synchronous gauge for $f(R)$ cosmology and study the impact of $f(R)$ gravity on the large scale structure formation. In section IV, we present the fittings results by confronting our model with the latest observations. In section V, we summarize and conclude this work.

II. THE BACKGROUND DYNAMICS

We work with the 4-dimensional action in the $f(R)$ gravity[13]

$$S = \frac{1}{2\kappa^2}\int d^4x\sqrt{-g}f(R) + \int d^4x\mathcal{L}^{(m)} \quad , \tag{1}$$

where $\kappa^2 = 8\pi G$. We consider a homogeneous and isotropic background universe described by the Friedmann-Robertson- Walker(FRW) metric

$$ds^2 = a^2[-d\tau^2 + d\sigma^2] \quad , \tag{2}$$

where $d\sigma^2$ is the conformal space-like hypersurface with a constant curvature $R^{(3)}=6K$

$$d\sigma^2=\frac{dr^2}{1-Kr^2}+r^2(d\theta^2+sin^2\theta d\phi^2) \quad . \tag{3}$$

The dynamics of the Universe in the $f(R)$ gravity is described by[13]

$$\ddot{F}+2F\dot{H}-H\dot{F}-\frac{2K}{a^2}=-\kappa^2(\rho+p) \quad . \tag{4}$$

where $F=\frac{df(R)}{dR}$, the dot denotes the time derivative with respect to the cosmic time t and ρ is the total energy density of the matter which consists of the cold dark matter, baryon and radiation $\rho=\rho_c+\rho_b+\rho_r$. p is the total pressure in the Universe. If we convert the derivatives in Eq.(4) from the cosmic time t to $x=\ln a$, Eq.(4) can be recast into

$$\frac{d^2}{dx^2}F+(\frac{1}{2}\frac{d\ln E}{dx}-1)\frac{dF}{dx}+(\frac{d\ln E}{dx}-\frac{2K}{E}e^{-2x})F = \frac{\kappa^2}{3E}\frac{d\rho}{dx} \quad , \tag{5}$$

where

$$\begin{aligned} E &\equiv \frac{H^2}{H_0^2} \quad , \\ R &\equiv 3(\frac{dE}{dx}+4E)+6Ke^{-2x} \quad , \\ \frac{d\rho}{dx} &= -3(\rho+p) \quad . \end{aligned} \tag{6}$$

For convenience, the energy density ρ, K and the scalar curvature R in Eq.(5) are in the unit of H_0^2 and we also set $\kappa^2=1$ in our analysis. In order to get a viable $f(R)$ model with a reasonable expansion history of the Universe and without loss of generality, we can parameterize the quantity $E(x)$ in Eq.(5) as the standard model in Einstein's gravity with an effective dark energy equation of state(EoS) w[6][14]

$$E(x)=\Omega_r^0e^{-4x}+\Omega_m^0e^{-3x}+\Omega_k^0e^{-2x}+\Omega_d^0e^{-3\int_0^x(1+w)dx} \quad , \tag{7}$$

where

$$\begin{aligned} \Omega_k^0 &\equiv -K \quad , \\ \Omega_m^0 &\equiv \frac{\kappa^2\rho_m^0}{3H_0^2} \quad , \\ \Omega_d^0 &\equiv \frac{\kappa^2\rho_d^0}{3H_0^2} \quad , \\ \Omega_r^0 &\equiv \frac{\kappa^2\rho_r^0}{3H_0^2} \quad . \end{aligned} \tag{8}$$

ρ_d^0 indicates the energy density of the effective dark energy, $\rho_m^0=\rho_b^0+\rho_c^0$ is the energy density of non-relativistic matter and $\rho^0=\rho_m^0+\rho_r^0$ is the total energy density of the matter.

The $f(R)$ model can be constructed by specifying the background expansion history of the Universe in Eq.(5). Eq.(5) becomes a second order differential equation only with respect to $F(x)$. However, any designed background expansion history is meaningless in physics because we have less knowledge about the nature of the dark energy equation of stat w at present. Therefore, it is of great interest to investigate the $f(R)$ model that can reproduce the same background expansion history as that of the ΛCDM model $w=-1$. Therefore in this work, we only focus on this family of $f(R)$ models. We numerically find the solution of Eq.(5) in order to include the curved case $\Omega_k^0\neq 0$. We perform our numerical calculation starting from the deep matter dominated epoch($a_i\sim 0.03$). The curvature term K can be neglected at this epoch and the analytical solutions of Eq.(5) give rise to

$$\begin{aligned} F(x) &\sim 1+D_m(e^{3x})^{p_+} \quad , \\ \frac{dF(x)}{dx} &\sim 3D_mp_+(e^{3x})^{p_+} \quad , \end{aligned} \tag{9}$$

where $p_+=\frac{5+\sqrt{73}}{12}$ and D_m is a dimensionless constant. In our work, we use the parameter D_m to characterize the family of $f(R)$ models instead of using the parameter B_0 proposed in [6] because, as presented in the appendix VI B, D_m directly relates to the covariant parameter ϖ in the Lagrangian formalism for this kind of $f(R)$ models in the spatially flat Universe. Since the $f(R)$ model investigated in this work and ΛCDM model can only be distinguished in their perturbed space-time, in the next section, we turn to the cosmological perturbations theory.

III. COSMOLOGICAL PERTURBATIONS

The cosmological perturbation theory for the $f(R)$ gravity has been well studied in [15]. [15] has extensively presented the perturbation equations for a wide family of modified gravities. In the appendix, we summarize the self-consistent covariant perturbation equations in the $f(R)$ gravity.

In this work, we use the synchronous gauge which is defined by $\psi=0$ and $B=0$. The synchronous gauge is widely used in the Einstein-Boltzmann codes [16] [17] [18] [19] to calculate the temperature and polarization power spectra of the cosmic microwave background anisotropy.

The synchronous gauge is characterized by the following two parameters

$$\begin{aligned} \eta_T &= -(\phi+\frac{E}{6}) \quad , \\ h_L &= 6\phi \quad , \end{aligned} \tag{10}$$

where η_T refers to the conformal 3-space curvature perturbation

$$\delta R^{(3)}=6\delta K=-4(k^2-3K)\eta_T \quad . \tag{11}$$

The perturbed modified Einstein equations are given by

$$\begin{aligned}
-\frac{1}{2}\kappa^2 a^2 \delta\rho &= -(\frac{1}{2}F\mathcal{H}+\frac{1}{4}F')h'_L-\frac{3}{2}\mathcal{H}\delta F'-\frac{1}{2}\delta F k^2 \\
&+ F\eta_T(k^2-3K)+\frac{3}{2}\mathcal{H}'\delta F \quad , \qquad (12)\\
\kappa^2 a^2 \delta p &= F[-\frac{2}{3}\mathcal{H}h'_L+\frac{2}{3}k^2\eta_T-\frac{1}{3}h''_L-2\eta_T K] \\
&+ \delta F[\mathcal{H}^2+\frac{a''}{a}-\frac{2}{3}k^2+2K]-\frac{1}{3}F'h'_L \\
&- \delta F''-\delta F'\mathcal{H} \quad , \qquad (13)\\
\alpha' &= -2\mathcal{H}\alpha+\eta_T-\frac{F'}{F}\alpha-\kappa^2 a^2\frac{p\Pi}{Fk^2}-\frac{\delta F}{F} \quad , \\
& \qquad (14)\\
\frac{k^2-3K}{k}F\eta'_T &= \frac{1}{2}\kappa^2 a^2 q+\frac{1}{2}k\delta F'-\frac{1}{2}k\mathcal{H}\delta F+\frac{Fh'_L K}{2k} \quad , \\
& \qquad (15)
\end{aligned}$$

where

$$\alpha \equiv \frac{(h_L+6\eta_T)'}{2k^2} \quad , \qquad (16)$$

and

$$\begin{aligned}
&\delta F''+2\mathcal{H}\delta F'+a^2(\frac{k^2}{a^2}+M^2)\delta F \\
&= \frac{\kappa^2 a^2}{3}(\delta\rho-3\delta p)-\frac{1}{2}F'h'_L \quad . \qquad (17)
\end{aligned}$$

The scalar curvature perturbation δR is given by

$$a^2\delta R = h''_T+3\mathcal{H}h'_T-4\eta_T k^2+12K\eta_T \quad . \qquad (18)$$

With the perturbation equations, we perform our numerical analysis based on the public available Einstein-Boltzmann code CAMB [18]. In the appendix, we summarize the details on how to modify the code. In Fig.1, we show the full spectrum of temperature anisotropy C_l^{TT} for a few representative values of D_m. The cosmological parameters used in the numerical process are taken from the WMAP 7-year best-fitted values for the ΛCDM model [2] $\Omega_b h^2 = 0.0227, \Omega_c h^2 = 0.112, n_s = 0.966, \Delta^2_{\mathcal{R}} = 2.42\times 10^{-9}, \Omega_\Lambda = 0.729, \tau = 0.085$. From Fig.1, we can see that the $f(R)$ gravity only affects the lowest multipoles in the power spectrum while not change the acoustic peaks. This observation is consistent with what found in[6]. The small l power is suppressed as the decreasing of the value of D_m. However, there is a turning around point, roughly about $D_m < -0.37 (B_0 > 1.5)$, after which there is a prominent enhancement in the power as further decreasing the value of D_m. Numerical results of our work agree with the already presented results within the parameterized framework of modified gravity[9][10].



FIG. 1: The angular powerspectrum of the temperature correlation

In Fig.2, we present the linear cold dark matter power spectrum at redshift zero $z = 0$. The $f(R)$ gravity changes not only the amplitude but also the shape of the power spectrum due to the scale k dependent growth history [6]. However, the $f(R)$ model investigated in our work should not change the position of the Baryon Acoustic Oscillation(BAO) peak in the two-point correlation function in the real space because the sound horizon is only determined by the background cosmological parameters[20][21] and the family of $f(R)$ models investigated in our work has the same background expansion. By doing the Fourier transformation

$$\xi(r) = \frac{1}{2\pi^2}\int dk k^2 P_L(k)\frac{\sin(kr)}{kr} \quad , \qquad (19)$$

in Fig 3, we present the two-point correlation function in the real space. We can see clearly that although the shape of the correlation function has changed a lot for a few representative values of D_m, the position of the BAO peak does not change under the Fourier transformation, which shows that our numerical results are consistent with the theoretical prediction.

In the spatially curved Universe, the curvature mainly shifts positions of acoustic peaks as shown in Fig.4. Although in the lowest multipoles, the impact of spatial curvature has similar behaviors as that of the modified gravity D_m, the high precision measurement of the acoustic peaks in the CMB temperature angular powerspectrum in combination with BAO and H_0[2] can put very tight constrains on the spatial curvature and the spatial curvature, therefore, could not have significant influence on the large scale structure in the $f(R)$ gravity.

Compared with the CMB temperature angular power spectrum C_l^{TT}(see Fig.1), there are no significant imprint of $f(R)$ gravity on the temperature polarization correlation spectrum and polarization auto-correlation spectrum C_l^{TE},C_l^{EE}(see Fig.5). After the qualitative analysis of the impact of $f(R)$ model on the large scale structure, in the next section, we will present cosmological constrains on the $f(R)$ model from latest observations.



FIG. 2: The linear cold dark matter powerspectrum.



FIG. 4: The impact of the curvature Ω_k on the angular powerspectrum of the temperature correlation



FIG. 3: The two-point correlation function in the real space



FIG. 5: The TE and EE angular powerspectrum.

IV. CONSTRAINS FROM OBSERVATIONS

The parameter space of our model is

$$P = (\Omega_b h^2, \Omega_m h^2, \theta_A, \Omega_k, \ln[10^{10}A_s], n_s, \tau, D_m) \quad , \quad (20)$$

where θ_A is the angular size of the acoustic horizon, and A_s is amplitude of the primordial curvature perturbation. To avoid the instabilities in the high curvature region, it requires that $D_m < 0$ [22]. Priors for cosmological parameters are listed in table I. For the CMB data, we use the seven-year WMAP(WMAP 7) CMB temperature and polarization power spectra [2]. We also take the measurement of the power spectrum of LRG from SDSS DR7 [23]. However, we limit our considerations of the matter power spectrum in a relatively linear scale and we only take the samples within bins for $k < 0.1h/\mathrm{Mpc}$. We add the supernovae data [24] and the present day value of the Hubble constant from the measurement of Hubble Space Telescope(HST) $H_0 = 74.2 \pm 3.6\mathrm{kms}^{-1}\mathrm{Mpc}^{-1}$ [25] in order to put tighter constrains on the background cosmological parameters. The global fitting results using above data sets are listed in Table II and III. In the flat Universe, we find the constrains on the parameter D_m as $|D_m| < 0.709(95\%\mathrm{CL})$ and $|D_m| < 0.711(95\%\mathrm{CL})$ in the curved Universe. These results are equivalent to $B_0 < 3.86(95\%\mathrm{CL})$ and $B_0 < 3.88(95\%\mathrm{CL})$ if we use the parameter B_0. The result is consistent with [11]. Although the $f(R)$ gravity has significant impact

TABLE I: Priors for cosmological parameters

$0.005 < \Omega_b h^2 < 0.1$
$0.01 < \Omega_c h^2 < 0.99$
$0.5 < \theta < 10$
$-0.2 < \Omega_k < 0.2$
$0.01 < \tau < 0.8$
$0.5 < n_s < 1.5$
$2.7 < \ln[10^{10}\mathrm{As}] < 4.0$
$-1.2 < D_m < 0$

on the shape of the matter power spectrum(see Fig. 2), the SDSS LRG matter spectrum can not put very tight constrains on D_m because there are degeneracies between D_m and other cosmological parameters[11]. When $D_m \sim -0.6(B_0 \sim 3)$, the CMB temperature angular power spectrum C_l^{TT} of the $f(R)$ model goes back similar to that of ΛCDM model. Therefore, there are two peaks in the likelihood distribution of D_m which are clearly shown in black lines in Fig.6 and Fig.7. Clearly, the combination of the data set CMB+SN+HST+MPK can not put very tight constrains on the $f(R)$ models. We need to add additional data set.

As pointed out in [11], the $f(R)$ gravity can produce the anti-correlation in the Galaxy-ISW angular power spectrum. However, the measurement of the Galaxy-ISW correlation favors the positive correlation. Therefore, the Galaxy-ISW correlation data set can put very tight constrains on the $f(R)$ model. We consider the cross correlation

$$C_l^{gI} = \frac{2}{\pi}\int k^2 dk P_m(k,0) W_l^g(k) W_l^I(k) \quad , \qquad (21)$$

where the window functions $W_l^I(k)$ and $W_l^g(k)$ are given by

$$\begin{aligned} W_l^I(k) &= -T_0 \int dz \frac{d}{dz}(\Psi - \Phi) j_l[k\chi(z)] \quad , \\ W_l^g(k) &= \int dz b(z)\Pi(z)D(z) j_l[k\chi(z)] \quad , \end{aligned} \qquad (22)$$

where $\chi(z)$ is the looking back time, $j_l(x)$ is the spherical bessel function and T_0 is the temperature of the CMB.

In order to improve the performance in the numerical process, we use the Limber approximation

$$\frac{2}{\pi}\int k^2 dk j_l[k\chi] j_l[k\chi'] \approx \frac{\delta(\chi - \chi')}{\chi^2} \quad , \qquad (23)$$

and Eq.(21) reduces to

$$C_l^{gI} \sim T_0 \int dz \frac{b(z)\Pi(z)}{\chi^2} P_{gI}(\frac{l+1/2}{\chi}, z) \quad , \qquad (24)$$

where

$$P_{gI}(k,z) = \frac{2\pi^2}{k^3} P_\chi(k) \delta_m(k,z) \frac{d}{d\tau}(\Psi - \Phi) \quad , \qquad (25)$$

and

$$P_\chi(k) = A_s(\frac{k}{k_*})^{n_s - 1} \quad . \qquad (26)$$

In this work, we adopt the Galaxy-ISW correlation data from [26] and use the public available ISWWLL code[26] to calculate the likelihood. We have turned off the contribution of weak lensing in the original code. The results of the joint likelihood analysis with the data sets from CMB+SN+HST+MPK+gISW are shown in Table II and III. The marginalized $1D$ and $2D$ likelihoods for interested parameters are shown in red lines in Fig.6 and Fig.7. The Galaxy-ISW correlation data set has improved significantly the constrains on the parameter of D_m up to $|D_m| < 0.109 (B_0 < 0.376)$(95%CL) in the flat Universe and $|D_m| < 0.131 (B_0 < 0.459)$(95%CL) in the curved Universe. The result in the flat Universe are in good agreement with the work done by [10]and [12]. The best fitted point for the curvature is slightly positive $\Omega_k = 0.0063^{+0.0061}_{-0.0061}$ suggesting that the Universe is nearly flat but open ($K < 0$), which is different from the results that the Universe is nearly flat but closed($K > 0$) in the non-flat ΛCDM model $\Omega_k = -0.0023^{+0.0054}_{-0.0056}$[2]. In the $f(R)$ gravity, our Universe may have a completely different expansion scenario in the far future, if the Universe is not exactly spatially flat.

V. CONCLUSIONS

In this work, we have studied the impact of the specific family of $f(R)$ models that can reproduce the same background expansion history as that of ΛCDM model on the cosmic microwave background and the large scale structure using the full set of covariant cosmological perturbation equations. Based upon the covariant perturbation equations, we have modified the public available Einstein-Boltzmann code CAMB and CosmoMC [27] to conduct the Markov chain Monte Carlo analysis on the parameter space and confront our $f(R)$ model with the latest observations. In the flat Universe, our results are in a good agreement with previous independent work within the parameterized framework of modified gravity. We also have extended our analysis to the non-flat Universe. From the fitting results, we find that the Universe is nearly flat but open at the best-fitted point in the $f(R)$ gravity which is different from the result that

TABLE II: Fitting results for the flat model $\Omega_k = 0$

Parameters	CMB+SN+HST+MPK	CMB+SN+HST+MPK+gISW
$\Omega_b h^2$	$0.02257^{+0.00053}_{-0.00053}$	$0.02257^{+0.00053}_{-0.00053}$
$\Omega_c h^2$	$0.1067^{+0.0040}_{-0.0040}$	$0.1055^{+0.0041}_{-0.0041}$
θ	$1.0396^{+0.0026}_{-0.0026}$	$1.0395^{+0.0027}_{-0.0027}$
τ	$0.091^{+0.015}_{-0.015}$	$0.091^{+0.015}_{-0.015}$
n_s	$0.966^{+0.013}_{-0.013}$	$0.968^{+0.013}_{-0.013}$
$\ln[10^{10}\mathrm{As}]$	$3.065^{+0.035}_{-0.035}$	$3.060^{+0.036}_{-0.036}$
$\lvert D_m \rvert$	< 0.709(95%CL)	< 0.109(95%CL)

TABLE III: Fitting results for the non-flat model $\Omega_k \neq 0$

Parameters	CMB+SN+HST+MPK	CMB+SN+HST+MPK+gISW
$\Omega_b h^2$	$0.02240^{+0.00055}_{-0.00055}$	$0.02246^{+0.00054}_{-0.00054}$
$\Omega_c h^2$	$0.1102^{+0.0052}_{-0.0052}$	$0.1084^{+0.0050}_{-0.0050}$
θ	$1.0390^{+0.0027}_{-0.0027}$	$1.0392^{+0.0025}_{-0.0025}$
τ	$0.088^{+0.014}_{-0.014}$	$0.089^{+0.015}_{-0.015}$
Ω_k	$0.0063^{+0.0061}_{-0.0061}$	$0.0051^{+0.0058}_{-0.0058}$
n_s	$0.962^{+0.014}_{-0.014}$	$0.964^{+0.014}_{-0.014}$
$\ln[10^{10}\mathrm{As}]$	$3.070^{+0.035}_{-0.035}$	$3.066^{+0.035}_{-0.035}$
$\lvert D_m \rvert$	< 0.711(95%CL)	< 0.131(95%CL)



FIG. 6: Marginalized posterior distribution and 2-D contour plots for $f(R)$ parameters in the flat Universe $\Omega_k = 0$. CMB+SN+HST+MPK(black lines), CMB+SN+HST+MPK+gISW(red lines)



FIG. 7: Marginalized posterior distribution and 2-D contour plots for $f(R)$ parameters in the non-flat Universe $\Omega_k \neq 0$. CMB+SN+HST+MPK(black lines), CMB+SN+HST+MPK+gISW(red lines)

the Universe is nearly flat but closed in the case of the ΛCDM model [2]. The modified gravity may change the final fate of our Universe in the future expansion, if the Universe is not exactly spatially flat.

Acknowledgment: J.H.He would like to thank B. R. Granett and L. Guzzo for helpful discussions. J.H.He acknowledges the Financial support of MIUR through PRIN 2008 and ASI through contract Euclid-NIS I/039/10/0.

VI. APPENDIX

A. Modifying CAMB

Our work is based on the public available Einstein-Boltzmann code CAMB [18]. CAMB propagates the covariant equations rest upon the CDM frame, which is equivalent to the synchronous gauge. Here we summarize the cosmological perturbation equations with the same conventions as that used in CAMB. In CAMB, the curvature perturbations are characterized by $\mathcal{Z}$ and σ

$$\begin{aligned} \mathcal{Z} &= \frac{h'_L}{2k} \quad , \\ \sigma &= k\alpha \quad , \end{aligned}$$

where

$$\eta'_T = \frac{k}{3}(\sigma - \mathcal{Z}) \quad . \tag{27}$$

The perturbed modified Einstein equations can be written as

$$
\begin{aligned}
(F\mathcal{H}+\frac{1}{2}F')k\mathcal{Z} &= \frac{\kappa^2}{2}a^2\delta\rho + Fk^2\eta_T\beta_2 - \frac{3}{2}\mathcal{H}\delta F' \\
&- \frac{1}{2}\delta F k^2 + \frac{3}{2}\mathcal{H}'\delta F \quad , \qquad (28) \\
\frac{k^2}{3}F(\beta_2\sigma - \mathcal{Z}) &= \frac{\kappa^2}{2}a^2 q + \frac{1}{2}k\delta F' \\
&- \frac{1}{2}k\mathcal{H}\delta F \quad , \qquad (29) \\
\sigma' + 2\mathcal{H}\sigma + \frac{F'}{F}\sigma &= k\eta_T - \kappa^2 a^2\frac{p\Pi}{Fk} - k\frac{\delta F}{F} \quad , \qquad (30) \\
\mathcal{Z}' + (\frac{1}{2}\frac{F'}{F}+\mathcal{H})\mathcal{Z} &= (-k\beta_2 + \frac{k}{2} + \frac{3\mathcal{H}^2}{k})\frac{\delta F}{F} \\
&- \frac{\kappa^2 a^2}{2kF}(\delta\rho + 3\delta p) - \frac{3}{2}\frac{\delta F''}{kF} \quad ,(31)
\end{aligned}
$$

where β_2 is the curvature factor

$$
\beta_2 = \frac{k^2-3K}{k^2} \quad . \qquad (32)
$$

The propagation of the perturbed field δF is given by

$$
\begin{aligned}
&\delta F'' + 2\mathcal{H}\delta F' + a^2(\frac{k^2}{a^2} + M^2)\delta F \\
&= \frac{\kappa^2 a^2}{3}(\delta\rho - 3\delta p) - kF'\mathcal{Z} \quad , \qquad (33)
\end{aligned}
$$

and the perturbation of the scalar curvature δR is given by

$$
a^2\delta R = 2kZ' + 6kZ\mathcal{H} - 4k^2\beta_2\eta_T \quad . \qquad (34)
$$

We have replaced the original perturbation equations with the above set of equations in the original CAMB code. Another important part we need to modify is the source term of the CMB temperature anisotropy [28][29]

$$
\begin{aligned}
&S_T(\tau,k) \\
&= e^{-\varepsilon}(\alpha'' + \eta_T') \\
&+ g(\Delta_{T0} + 2\alpha' + \frac{v_b'}{k} + \frac{\zeta}{12\sqrt{\beta_2}} + \frac{\zeta''}{4k^2\sqrt{\beta_2}}) \\
&+ g'(\alpha + \frac{v_b}{k} + \frac{\zeta'}{2k^2\sqrt{\beta_2}}) + \frac{1}{4}\frac{g''\zeta}{k^2\sqrt{\beta_2}} \qquad (35) \\
&= e^{-\varepsilon}(\frac{\sigma''}{k} + \frac{k\sigma}{3} - \frac{k\mathcal{Z}}{3}) \\
&+ g(\Delta_{T0} + 2\frac{\sigma'}{k} + \frac{v_b'}{k} + \frac{\zeta}{12\sqrt{\beta_2}} + \frac{\zeta''}{4k^2\sqrt{\beta_2}}) \\
&+ g'(\frac{\sigma}{k} + \frac{v_b}{k} + \frac{\zeta'}{2k^2\sqrt{\beta_2}}) + \frac{1}{4}\frac{g''\zeta}{k^2\sqrt{\beta_2}} \quad ,
\end{aligned}
$$

where $g = -\dot{\varepsilon}e^{-\varepsilon} = an_e\sigma_T e^{-\varepsilon}$ is the visibility function and ε is the optical depth. ζ is given by

$$
\zeta = (\frac{3}{4}I_2 + \frac{9}{2}E_2) \quad , \qquad (36)
$$

where I_2, E_2 indicate the quadrupole of the photon intensity and the E-like polarization respectively [29]. In the original code, σ and $\mathcal{Z}$ are calculated from the perturbation equations in the standard Einstein's gravity. In our work, σ and $\mathcal{Z}$ are calculated from perturbation equations in the $f(R)$ gravity.

B. Lagrangian formalism

In the spatially flat Universe filled with dust matter, the analytical solution of Eq.(5) reads

$$
\begin{aligned}
F(x) - 1 &= C_m(e^{3x})^{p_-}\,{}_2F_1[q_-,p_-;r_-;-e^{3x}\frac{\Omega_d^0}{\Omega_m^0}] \\
&+ D_m(e^{3x})^{p_+}\,{}_2F_1[q_+,p_+;r_+;-e^{3x}\frac{\Omega_d^0}{\Omega_m^0}] \quad , \qquad (37)
\end{aligned}
$$

where the indexes are

$$
\begin{aligned}
q_+ &= \frac{1+\sqrt{73}}{12} \quad , \\
q_- &= \frac{1-\sqrt{73}}{12} \quad , \\
r_+ &= 1+\frac{\sqrt{73}}{6} \quad , \\
r_- &= 1-\frac{\sqrt{73}}{6} \quad , \\
p_+ &= \frac{5+\sqrt{73}}{12} \quad , \\
p_- &= \frac{5-\sqrt{73}}{12} \quad . \qquad (38)
\end{aligned}
$$

The viable $f(R)$ model should be in a "chameleon" type [30][31] where in the past of the Universe, the model should go back to the standard model

$$
\lim_{x\to-\infty} F(x) = 1 \quad . \qquad (39)
$$

C_m, therefore, should vanish in the solution $C_m = 0$.

The Lagrangian formalism for the corresponding $f(R)$ model is given by

$$
\begin{aligned}
f(R) &= R - 2\Lambda \\
&- \varpi\left(\frac{\Lambda}{R-4\Lambda}\right)^{p_+-1}\,{}_2F_1\left[q_+,p_+-1;r_+;-\frac{\Lambda}{R-4\Lambda}\right] \quad , \qquad (40)
\end{aligned}
$$

where Λ, ϖ are constant parameters.

$$
\varpi = D_m(R_0-4\Lambda)^{p_+}/(p_+-1)/\Lambda^{p_+-1} \quad . \qquad (41)
$$

When $\varpi = 0$, Λ is just the cosmological constant. The above expression is a well-defined real function on the real axes in the physical range $R > 4\Lambda$, which is different from the results given by [32]. Moreover, the Lagrangian formalism does not have the future singularity because $f(R)$ is finite at the point of $R = 4\Lambda$.

$$\begin{aligned} &\lim_{R\to 4\Lambda} f_2(R) = 2\Lambda - \\ &\frac{\varpi 4(-511+79\sqrt{73})\Gamma(2/3)\Gamma(-r_-)}{(-5+\sqrt{73})(-1+\sqrt{73})(7+\sqrt{73})\Gamma(-p_-)\Gamma(q_+)} \quad , \\ &\approx 2\Lambda - 1.256\varpi \end{aligned} \tag{42}$$

where Γ is the Euler Gamma function.

When $R < 4\Lambda$ the expression of Eq.(40) becomes complex, which is clearly unphysical.

C. Cosmological perturbation in f(R) gravity

The perturbed line element for the scalar perturbation can be written as[33]

$$\begin{aligned} ds^2 &= a^2[-(1+2\psi Y^{(s)})d\tau^2 + 2BY_i^{(s)}d\tau dx^i \\ &+ (1+2\phi Y^{(s)})\gamma_{ij}dx^i dx^j + EY_{ij}^{(s)}dx^i dx^j] \quad . \end{aligned} \tag{43}$$

γ_{ij} in the spherical coordinate can be written as

$$[\gamma_{ij}] = \begin{pmatrix} \frac{1}{1-Kr^2} & 0 & 0 \\ 0 & r^2 & 0 \\ 0 & 0 & r^2 sin^2\theta \end{pmatrix} \quad , \tag{44}$$

where we also include the spatially curved case. Perturbation equations in the spatially flat Universe can also be found in [34]. Y,Y_j and Y_{ij} are the scalar harmonic function, which are defined by

$$\begin{aligned} &(\Delta + k^2)Y^{(s)} = 0 \quad , \\ &Y_j^{(s)} \equiv -\frac{1}{k}Y_{|j}^{(s)} \quad , \\ &Y_{ij}^{(s)} \equiv \frac{1}{k^2}Y_{|ij}^{(s)} + \frac{1}{3}\gamma_{ij}Y^{(s)} \quad . \end{aligned} \tag{45}$$

The perturbed modified Einstein equations are given by

$$\begin{aligned} -\frac{\kappa^2}{2}\delta\rho a^2 &= F[-k^2\phi + 3\mathcal{H}(\mathcal{H}\psi - \phi') \\ &+ k\mathcal{H}B + K(3\phi + \frac{1}{2}E) - \frac{1}{6}k^2E] \\ &+ F'(\frac{k}{2}B + 3\mathcal{H}\psi - \frac{3}{2}\phi') - \frac{3}{2}\mathcal{H}\delta F' \\ &- \delta F(\frac{3}{2}\mathcal{H}^2 - \frac{3}{2}\frac{a''}{a} + \frac{k^2}{2}) \quad . \end{aligned} \tag{46}$$

$$\begin{aligned} \kappa^2 a^2\delta p &= F[-\frac{2}{3}k^2\psi - \frac{2}{3}k^2\phi - \frac{1}{9}k^2E \\ &+ 4\frac{a''}{a}\psi + \frac{2}{3}kB' + \frac{4}{3}k\mathcal{H}B - 2\mathcal{H}^2\psi \\ &+ 2\mathcal{H}\psi' - 4\mathcal{H}\phi' - 2\phi'' + K(2\phi + \frac{1}{3}E)] \\ &+ F'[\frac{2}{3}kB + \psi' + 2\mathcal{H}\psi - 2\phi'] \\ &+ \delta F[\mathcal{H}^2 + \frac{a''}{a} - \frac{2}{3}k^2 + 2K] \\ &- \delta F'' - \delta F'\mathcal{H} + 2\psi F'' \quad . \end{aligned} \tag{47}$$

$$\begin{aligned} \kappa^2 a^2 p\Pi &= F[-k^2\psi - k^2\phi + 2k\mathcal{H}B + \mathcal{H}E' \\ &- \frac{1}{6}k^2E + \frac{1}{2}E'' + kB'] \\ &+ F'(\frac{1}{2}E' + kB) - k^2\delta F \quad . \end{aligned} \tag{48}$$

$$\begin{aligned} -\frac{1}{2}\kappa^2 a^2 q &= F[k\phi' - k\mathcal{H}\psi + 2\mathcal{H}^2B \\ &- \frac{a''}{a}B + \frac{1}{6}kE' - \frac{1}{2}E'\frac{K}{k}] \\ &+ F'[\mathcal{H}B - \frac{1}{2}k\psi] + \frac{1}{2}k\delta F' \\ &- \frac{1}{2}k\mathcal{H}\delta F - \frac{1}{2}BF'' \quad , \end{aligned} \tag{49}$$

where $q = (\rho+p)v$. The above perturbation equations are self-consistent covariant equations. We can show that under the infinitesimal coordinate transformation, namely, Eq.(50), the perturbation equations could keep the same form.

$$\begin{aligned} \hat{\psi}Y^{(s)} &= (\psi - \xi^{0\prime} - \mathcal{H}\xi^0)Y^{(s)} \quad , \\ \hat{\phi}Y^{(s)} &= (\phi - \frac{1}{3}k\beta - \mathcal{H}\xi^0)Y^{(s)} \quad , \\ \hat{B}Y_i^{(s)} &= (B - k\xi^0 - \beta')Y_i^{(s)} \quad , \\ \hat{E}Y_{ij}^{(s)} &= (E + 2k\beta)Y_{ij}^{(s)} \quad . \end{aligned} \tag{50}$$

The perturbation for the scalar field δF satisfies

$$\begin{aligned} \delta F &= f_{RR}\delta R \quad , \\ f_{RR} &= \frac{\partial F}{\partial R} \quad , \end{aligned}$$

where the perturbation of scalar curvature δR is given by

$$\begin{aligned} a^2\delta R = -12\psi\frac{a''}{a} - 6\mathcal{H}\psi' + 4\phi k^2 - 2KE \\ + 2\psi k^2 + 18\mathcal{H}\phi' + \frac{2}{3}Ek^2 + 6\phi'' \\ - 6B\mathcal{H}k - 12\phi K - 2B'k \quad . \end{aligned} \tag{51}$$

Inserting Eq.(51) into Eq.(47) to eliminate ϕ'' and using Eq.(46) to eliminate $k^2\phi$, the equation governing the behavior of δF gives rise to

$$\begin{aligned}&\delta F'' + 2\mathcal{H}\delta F' + a^2(\frac{k^2}{a^2} + M^2)\delta F \\ &= \frac{\kappa^2 a^2}{3}(\delta\rho_m - 3\delta p_m) + 2\psi F'' \\ &+ F'(4\mathcal{H}\psi - 3\phi' + \psi' + kB) \quad ,\end{aligned} \tag{52}$$

where

$$M^2 = \frac{1}{3}(\frac{F}{f_{RR}} - R) \quad , \tag{53}$$

and

$$R = \frac{6a''}{a^3} + \frac{6K}{a^2} \quad . \tag{54}$$

---

[1] S. J. Perlmutter et al., Nature 391 (1998) 51; A. G. Riess et al., Astron. J. 116 (1998) 1009 ; S. J. Perlmutter et al., Astroph. J. 517 (1999) 565 ; J. L. Tonry et al., Astroph. J. 594 (2003) 1; A. G. Riess et al., Astroph. J. 607 (2005) 665 ; P. Astier et al., Astron. Astroph. 447 (2006) 31 ; A G. Riess et al., Astroph. J. 659 (2007) 98.

[2] E. Komatsu, et.al., ApJS 192 (2011) 18, arXiv:1001.4538.

[3] Ariel G. Sanchez, et.al. arXiv:1203.6616.

[4] Sean M. Carroll, LivingRev. Rel. 4 (2001) 1, astro-ph/0004075.

[5] P.G. Bergmann, Int. J. Theor. Phys. 1 (1968) 25;A. A. Starobinsky, Phys. Lett. B 91 (1980) 99; A.L. Erickcek, T.L. Smith, T.L., and M. Kamionkowski, Phys. Rev. D 74 (2006) 121501; V. Faraoni, Phys. Rev. D 74 (2006) 023529; S. Capozziello, and S. Tsujikawa, Phys. Rev. D 77 (2008) 107501 ;T. Chiba, T.L. Smith and A. L. Erickcek, Phys. Rev. D 75 (2007) 124014; I. Navarro, and K. Van Acoleyen, J. Cosmol. Astropart. Phys. 02 (2007) 022; G. J. Olmo, Phys. Rev. Lett. 95 (2005) 261102;G. J. Olmo, Phys. Rev. D 72 (2005) 083505; Amendola, L., Polarski, D., and Tsujikawa, S., Phys. Rev. Lett. 98 (2007) 131302, astro-ph/0603703; Luca Amendola, Radouane Gannouji, David Polarski, Shinji Tsujikawa, Phys.Rev.D 75 (2007) 083504, gr-qc/0612180; Luca Amendola, Phys.Rev.D 60 (1999) 043501, astro-ph/9904120.

[6] Yong-Seon Song, Wayne Hu, and Ignacy Sawicki, Phys.Rev. D 75 (2007) 044004 ,astro-ph/0610532;

[7] Jian-Hua He, Bin Wang, Elcio Abdalla, Phys. Rev. D 84 (2011) 123526, arXiv:1109.1730.

[8] W. Hu, I. Sawicki, Phys.Rev. D 76 (2007) 064004, arXiv:0705.1158;

[9] W. Hu, I. Sawicki, Phys. Rev. D 76 (2007) 104043, arXiv:0708.1190; W. Hu, Phys. Rev. D 77 (2008) 103524 , arXiv:0801.2433; W. Fang et al., Phys. Rev. D 78 (2008) 103509 , arXiv:0808.2208; W. Fang, W. Hu, A. Lewis, Phys. Rev. D 78 (2008) 087303 , arXiv:0808.3125.

[10] Alireza Hojjati, Levon Pogosian, Gong-Bo Zhao, JCAP 1108 (2011) 005, arXiv:1106.4543; Gong-Bo Zhao, Levon Pogosian, Alessandra Silvestri and Joel Zylberberg, Phys.Rev.D 79 (2009) 083513, arXiv:0809.3791

[11] Yong-Seon Song, Hiranya Peiris, Wayne Hu, Phys.Rev.D76 (2007)063517, arXiv:0706.2399

[12] Lucas Lombriser, Anze Slosar, Uros Seljak, Wayne Hu, arXiv:1003.3009

[13] Alessandra Silvestri, Mark Trodden, Rept.Prog.Phys.72 (2009) 096901, arXiv:0904.0024; Antonio De Felice, Shinji Tsujikawa, LivingRev. Rel. 13 (2010) 3 , arXiv:1002.4928; Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, Constantinos Skordis, Physics Reports 513 (2012) 1 ; Thomas P. Sotiriou, Valerio Faraoni, Rev. Mod. Phys. 82 (2010) 451, arXiv:0805.1726

[14] Levon Pogosian, Alessandra Silvestri, Phys.Rev.D77 (2008) 023503, arXiv:0709.0296

[15] Hwang, J.-C., Noh, H. Phys. Rev. D 65 (2001) 023512;Hwang, J.-C., and Noh, H. Phys. Rev. D, 71 (2005) 063536;

[16] U. Seljak and M. Zaldarriaga, Astrophys. J. 469, 437 (1996), astro-ph/9603033.

[17] Michael Doran, JCAP 0510 (2005) 011, astro-ph/0302138

[18] A. Lewis, A. Challinor and A. Lasenby, Astrophys. J.538 (2000) 473, arXiv:astro-ph/9911177.

[19] J. Lesgourgues, arXiv:1104.2932; D. Blas, J. Lesgourgues, T. Tram, JCAP 1107 (2011) 034, arXiv:1104.2933; J. Lesgourgues, arXiv:1104.2934; J. Lesgourgues, T. Tram, JCAP 1109 (2011) 032, arXiv:1104.2935; B. Audren, J. Lesgourgues, JCAP 1110 (2011) 037, arXiv:1106.2607

[20] Bruce A. Bassett, Rene Hlozek, Dark Energy, Ed. P. Ruiz-Lapuente (2010, ISBN-13: 9780521518888) ,arXiv:0910.5224

[21] Eisenstein, D. and White, M. Phys. Rev. D, 70 (2004) 103523 .

[22] Ignacy Sawicki, Wayne Hu, Phys. Rev. D 75 (2007) 127502, arXiv:astro-ph/0702278.

[23] B. Reid et al., Mon. Not. Roy. Astron. Soc. 404 (2010) 60.

[24] R. Amanullah, et al., Astrophys. J. 716 (2010) 712. arXiv:1004.1711.

[25] A. G. Riess et al., Astrophys. J. 699 (2009) 539, arXiv:0905.0695

[26] S. Ho, C. Hirata, N. Padmanabhan, U. Seljak and N. Bahcall, Phys. Rev, D78, (2008) 043519, arXiv:0801.0642; C. M. Hirata, S. Ho, N. Padmanabhan, U. Seljak and N. A. Bahcall, Phys. Rev. D 78, (2008) 043520, arXiv:0801.0644.

[27] A. Lewis and S. Bridle, Phys. Rev. D66 (2002) 103511,astro-ph/0205436

[28] M. Zaldarriaga, U. Seljak and E. Bertschinger, Astrophys. J. 494 (1998) 491, astro-ph/9704265;

[29] Anthony Challinor, Phys.Rev. D62 (2000) 043004, astro-ph/9911481.

[30] David F. Mota, John D. Barrow, Phys.Lett.B581:141-146,2004, astro-ph/0306047

[31] Khoury, J., and Weltman, A., Phys. Rev. D 69 (2004) 044026 ; Khoury, J., and Weltman, A., Phys. Rev. Lett.,

93 (2004) 171104.

[32] A. de la Cruz-Dombriz, A. Dobado. Phys.Rev. D74 (2006) 087501, gr-qc/0607118; P. K. S. Dunsby et al., Phys.Rev. D82 (2010) 023519, arXiv:1005.2205; S. Nojiri and S. D. Odintsov, Phys. Rev. D74 (2006) 086005, hep-th/0608008; Shin'ichi Nojiri, Sergei D. Odintsov J.Phys.A40 (2007) 6725, hep-th/0610164; S. Capozziello, S. Nojiri, S. D. Odintsov, A. Troisi, Phys. Lett. B 639(2006) 135, astro-ph/0604431.

[33] H. Kodama, M. Sasaki, Prog. Theor. Phys. Suppl. 78 (1984) 1.

[34] Rachel Bean, David Bernat, Levon Pogosian, Alessandra Silvestri, Mark Trodden Phys.Rev.D75 (2007) 064020, astro-ph/0611321